

# CONTROL III®

## DISINFECTANT GERMICIDE

**ACTIVE INGREDIENTS:**

n-alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chloride..... 10%

n-alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chloride ..... 10%

**INERT INGREDIENTS** ..... 80%

**TOTAL** ..... 100%

**ON INANIMATE HARD, NON-POUROUS, PRE-CLEANED ENVIRONMENTAL SURFACES EFFECTIVE AGAINST**

- Staphylococcus aureus • Vaccinia viruses
- Salmonella choleraesuis • Influenza A<sub>2</sub> virus
- Escherichia coli • Herpes simplex virus
- Staphylococcus aureus (MRSA) • HIV-1

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See side panel for additional precautionary statements.

This product is a concentrate and must be diluted before use.

**NET 16 FL. OZ. (1 PINT)**  
EPA Est. No. 40873 CA-01  
EPA Reg. No. 55364-3

**MARIL PRODUCTS, INC.**  
15421 Red Hill Ave., Suite D, Tustin, California 92780 U.S.A.  
**Manufactured for Maril Products, Inc.**  
**(714) 544-7711 / www.controlthree.com**

**USED FOR:**  
Disinfection of walls, equipment, heat sensitive instruments and tubing in operating rooms, recovery rooms and intensive care rooms.

**DIRECTIONS FOR USE:**  
It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high-level disinfection.

- General:**
1. Remove soil as required.
  2. Apply this product as directed.

**NON-CRITICAL INSTRUMENT DISINFECTION**  
Clean non-critical instruments with detergent and rinse with potable water before disinfection. Prepare use solution of 1 ounce per gallon of water. Immerse non-critical instruments for 10 minutes. Remove instruments, rinse thoroughly with sterile water and pat dry or allow to air dry. Active concentration is 1562 ppm.

**HIV-1 AND HOME CARE INSTRUCTIONS**

See Technical Bulletins for Home Care and Efficacy of Control III Products for the Control of the Human Immunodeficiency Virus Type 1 (AIDS Virus).\*

**LAUNDRY BACTERIOSTAT FOR COMMERCIAL, INDUSTRIAL, AND NON-MEDICAL INSTITUTIONAL LAUNDRY APPLICATIONS:**

This product provides the fabric with residual bacteriostatic activity against odor-causing gram-negative and gram-positive bacteria when this product is added to the final rinse at a rate of 1.2 ounces per 25 lbs. (4.8 ounces per 100 lbs.) of dry laundry. This is based on water being at full capacity, temperature of 20-38° C and 5 minute contact.

**HOSPITAL DISINFECTION**  
For disinfecting walls, non critical equipment and surfaces in operating rooms. Pre-clean surfaces with product prior to disinfection. Thoroughly wet pre-cleaned inanimate hard, non-porous environmental surfaces by applying a use solution of 1oz. per gallon of water with cloth, mop, sponge, or sprayer. Allow surfaces to remain wet for 10 minutes, then allow to air dry. Prepare a fresh solution for each use. Active concentration is 1562 ppm.

**PRECAUTIONARY STATEMENTS:**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER:**  
Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield, protective clothing and rubber gloves when handling. May be harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling and before eating, drinking or smoking tobacco. Remove and wash contaminated clothing.

**STORAGE AND DISPOSAL:**  
Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:**  
Store in original container in a locked storage area inaccessible to children.  
**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.


**CONTAINER HANDLING:**  
Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Store rinsate for later disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or wrap container and discard in the trash.

**FIRST AID**  
**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**If on skin or clothing:** Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.  
**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**Note to Physician:**  
Probable mucosal damage may contraindicate the use of gastric lavage.

**HOT LINE NUMBER**  
For emergency medical advice, call your local poison control center (1-800-222-1222) or doctor. Have the product container or label with you when seeking medical advice or treatment.



7 22540 00002 8